



G. 89



1997

Budker Institute of Nuclear Physics
SB RAS

A.G. Grozin

RECENT PROGRESS ON HQET LAGRANGIAN

Budker INP 97-90



Novosibirsk

v

RECENT PROGRESS ON HQET LAGRANGIAN

A. G. Grozin

Budker Institute of Nuclear Physics
630090, Novosibirsk, Russian Federation

Abstract

HQET lagrangian up to $1/m^3$ terms is discussed. Consequences of reparameterization invariance are considered. Results for the chromomagnetic interaction coefficient at two loops, and in all orders in the large- β_1 approximation, are presented.

©Budker Institute of Nuclear Physics

1. HQET lagrangian

QCD problems with a single heavy quark staying approximately at rest can be conveniently treated in the heavy quark effective field theory (HQET) (see [1] for review and references). We shift the energy zero level: $E = m + \omega$, and consider the region where residual energies ω and momenta \vec{p} are not large: $\omega \sim |\vec{p}| \sim \Lambda \ll m$. The effective field theory is constructed to reproduce QCD on-shell scattering amplitudes expanded to some order $(\Lambda/m)^n$. This is achieved by writing down the most general effective Lagrangian consistent with the required symmetries, and tuning the coefficients to reproduce QCD on-shell amplitudes. Terms with $D_0 Q$ can be eliminated by field redefinitions.

The most general lagrangian up to $1/m^3$ is [2]–[6]

$$\begin{aligned}
 L = & Q^\dagger i D_0 Q \\
 & + \frac{C_k}{2m} Q^\dagger \vec{D}^2 Q + \frac{C_m}{2m} Q^\dagger \vec{B} \cdot \vec{\sigma} Q \\
 & + \frac{iC_s}{8m^2} Q^\dagger (\vec{D} \times \vec{E} - \vec{E} \times \vec{D}) \cdot \vec{\sigma} Q + \frac{C_d}{8m^2} Q^\dagger [\vec{D} \cdot \vec{E}] Q \\
 & + \frac{C_{k2}}{8m^3} Q^\dagger \vec{D}^4 Q + \frac{C_{w1}}{8m^3} Q^\dagger \{ \vec{D}^2, \vec{B} \cdot \vec{\sigma} \} Q - \frac{C_{w2}}{4m^3} Q^\dagger D^i \vec{B} \cdot \vec{\sigma} D^i Q \\
 & + \frac{C_{p'p}}{8m^3} Q^\dagger (\vec{D} \vec{B} \cdot \vec{D} + \vec{D} \cdot \vec{B} \vec{D}) \cdot \vec{\sigma} Q \\
 & + \frac{iC_M}{8m^3} Q^\dagger (\vec{D} \cdot [\vec{D} \times \vec{B}] + [\vec{D} \times \vec{B}] \cdot \vec{D}) Q \\
 & + \frac{C_{a1}}{8m^3} Q^\dagger (\vec{B}^2 - \vec{E}^2) Q - \frac{C_{a2}}{16m^3} Q^\dagger \vec{E}^2 Q \\
 & + \frac{C_{a3}}{8m^3} Q^\dagger \text{Tr}(\vec{B}^2 - \vec{E}^2) Q - \frac{C_{a4}}{16m^3} Q^\dagger \text{Tr} \vec{E}^2 Q \\
 & + \frac{iC_{b1}}{8m^3} Q^\dagger (\vec{B} \times \vec{B} - \vec{E} \times \vec{E}) \cdot \vec{\sigma} Q - \frac{iC_{b2}}{8m^3} Q^\dagger (\vec{E} \times \vec{E}) \cdot \vec{\sigma} Q + \dots
 \end{aligned} \tag{1}$$

where Q is 2-component heavy-quark field. Here heavy-light contact interactions are omitted, as well as operators involving only light fields.

HQET can be rewritten in relativistic notations. Momenta of all states are decomposed as $p = mv + k$ where residual momenta $k \sim \Lambda$. The heavy-quark field is now Dirac spinor obeying $\not{v}Q_v = Q_v$. The lagrangian is

$$L_v = \bar{Q}_v i v \cdot D Q_v - \frac{C_k}{2m} \bar{Q}_v D_\perp^2 Q_v - \frac{C_m}{4m} \bar{Q}_v G_{\mu\nu} \sigma^{\mu\nu} Q_v + \frac{iC_s}{8m^2} \bar{Q}_v \{D_\perp^\mu, G^{\lambda\nu}\} v_\lambda \sigma_{\mu\nu} Q_v - \frac{C_d}{8m^2} \bar{Q}_v v^\mu [D_\perp^\nu G_{\mu\nu}] Q_v + \dots \quad (2)$$

where $D_\perp = D - v(vD)$. The velocity v may be changed by an amount $\delta v \lesssim \Lambda/m$ without spoiling the applicability of HQET and changing its predictions. This reparameterization invariance relates coefficients of varying degrees in $1/m$ [7]–[13].

At the tree level, there are easier ways to find the coefficients C_i than QCD/HQET matching: Foldy-Wouthuysen transformation [14,15], or using equations of motion [5] (or integrating out lower components [16,17]) followed by a field redefinition. The result is

$$C_k = C_m = C_d = C_s = C_{k2} = C_{w1} = C_{a1} = C_{b1} = 1, \quad (3)$$

$$C_{w2} = C_{p'p} = C_M = C_{a2} = C_{a3} = C_{a4} = C_{b2} = 0.$$

However, these algebraic methods don't generalize to higher loops.

At $1/m$ level, the kinetic coefficient $C_k = 1$ due to the reparameterization invariance [7]. One-loop matching for the chromomagnetic coefficient C_m was done in [3]; two-loop anomalous dimension of the chromomagnetic operator in HQET was obtained in [18,19], and two-loop matching was done in [19]; in [20], all orders of perturbation theory for C_m were summed at large β_1 .

At $1/m^2$ level, the spin-orbit coefficient $C_s = 2C_m - 1$ due to the reparameterization invariance [21]–[24]. The Darwin term reduces to a contact interaction. One-loop matching for the heavy-light contact interactions was done in [24]. The one-loop anomalous dimension matrix of dimension 6 terms in the HQET lagrangian was obtained in [15], [22]–[25].

At $1/m^3$ level, one-loop matching was done in [6] for the terms involving the heavy-quark fields twice and the gluon field once. The one-loop renormalization of dimension 7 terms in the HQET lagrangian was recently considered [26].

2. Matching quark-quark vertex

Renormalized QCD on-shell quark-quark proper vertex

$$-\bar{u}(\not{p} - m)u \quad (4)$$

gets no correction in the on-shell renormalization scheme. QCD spinors are related to HQET spinors by the Foldy-Wouthuysen transformation

$$u = \left(1 + \frac{\not{k}}{2m} + \frac{k^2}{4m^2} + \dots\right) u_v, \quad \not{v}u_v = u_v. \quad (5)$$

Expressing QCD proper vertex via HQET spinors, we obtain

$$\bar{u}_v \frac{\not{k}^2}{2m} u_v + \dots \quad (6)$$

Let's denote the sum of bare 1-particle-irreducible self-energy diagrams of the heavy quark in HQET at $1/m^0$ as $-i\frac{1+\not{v}}{2}\Sigma(\omega)$, $\omega = kv$. At the $1/m$ level, self-energy diagrams with a single chromomagnetic vertex vanish. Let the sum of bare diagrams with a single kinetic vertex be $-i\frac{C_k}{2m}\frac{1+\not{v}}{2}\Sigma_k(\omega, k_\perp^2)$. Consider variation of Σ at $v \rightarrow v + \delta v$ for an infinitesimal δv ($v\delta v = 0$). All factors $\frac{1+\not{v}}{2}$ can be combined into a single one, and the variation $\delta\phi$ in it provides the variation of the γ -matrix structure in front of Σ . There are two sources of the variation of Σ . Terms from the expansion of denominators of the propagators produce insertions $ik\delta v$. Terms from the vertices produce $igt^a\delta v^\mu$. Now consider variation of Σ_k at $k_\perp \rightarrow k_\perp + \delta k_\perp$ for an infinitesimal δk_\perp . Quark-quark kinetic vertices produce $i\frac{C_k}{m}k\delta k_\perp$; quark-quark-gluon kinetic vertices produce $i\frac{C_k}{m}gt^a\delta k_\perp^\mu$; two-gluon vertices produce nothing. Therefore,

$$\frac{\partial\Sigma_k}{\partial k_\perp^\mu} = 2\frac{\partial\Sigma}{\partial v^\mu}. \quad (7)$$

This is the Ward identity of the reparameterization invariance first derived in [10]. Taking into account $\frac{\partial\Sigma_k}{\partial k_\perp^\mu} = 2\frac{\partial\Sigma_k}{\partial k_\perp^2}k_\perp^\mu$ and $\frac{\partial\Sigma}{\partial v^\mu} = \frac{d\Sigma}{d\omega}k_\perp^\mu$, we obtain

$$\frac{\partial\Sigma_k}{\partial k_\perp^2} = \frac{d\Sigma}{d\omega}. \quad (8)$$

The right-hand side does not depend on k_\perp^2 , and hence

$$\Sigma_k(\omega, k_\perp^2) = \frac{d\Sigma(\omega)}{d\omega}k_\perp^2 + \Sigma_{k0}(\omega). \quad (9)$$

This result can also be understood in a more direct way. Only diagrams with a quark-quark kinetic vertex contain k_{\perp}^2 ; its coefficient is $i\frac{C_k}{2m}$. The sum of diagrams with a unit insertion is $-i\frac{d\Sigma}{d\omega}$. Note that diagrams with a quark-quark-gluon kinetic vertex vanish because there is no preferred transverse direction.

On the mass shell ($\omega = 0$), the renormalized HQET quark-quark proper vertex is $\frac{C_k}{2m} Z_Q \bar{u}_v [-k_{\perp}^2 + \Sigma_k(0, k_{\perp}^2)] u_v = -\frac{C_k}{2m} Z_Q [1 - \frac{d\Sigma}{d\omega}]_{\omega=0} k_{\perp}^2 \bar{u}_v u_v$. On the mass shell, only diagrams with finite-mass particles in loops contribute (e.g., c -quark loops in b -quark HQET) (Fig. 1). Taking into account $Z_Q^{-1} = 1 - \frac{d\Sigma}{d\omega}|_{\omega=0}$ and comparing with (6), we finally obtain

$$C_k(\mu) = 1. \quad (10)$$

This argument works for an arbitrary μ ; hence, the anomalous dimension of the kinetic-energy operator in HQET vanishes exactly. In a similar way, it is not difficult to prove that

$$C_{k2} = 1. \quad (11)$$

$$Z_Q \left[\text{tree} + \text{loop}_1 + \text{loop}_2 + \text{loop}_3 + \text{loop}_4 + \dots \right] = Z_Q \left[1 + \text{loop}_1 \right] C_k \frac{k^2}{2m}$$

$$Z_Q^{-1} = 1 - \frac{d\Sigma}{d\omega} = 1 + \text{loop}_1 + \dots$$

Figure 1: HQET quark-quark proper vertex on the mass shell

3. Matching quark-quark-gluon vertex

QCD on-shell proper vertex is characterized by 2 form factors:

$$\bar{u}(p') t_a^\mu \left(\varepsilon(q^2) \frac{(p+p')^\mu}{2m} + \mu(q^2) \frac{[\not{q}, \gamma^\mu]}{4m} \right) u(p), \quad (12)$$

$$\varepsilon(q^2) = 1 + \varepsilon' \frac{q^2}{m^2} + \dots, \quad \mu(q^2) = \mu + \mu' \frac{q^2}{m^2} + \dots$$

The total colour charge of a quark $\varepsilon(0) = 1$ due to the gauge invariance. Ward identities in the background field formalism [27] are shown in Fig. 2, where the large dot means convolution with the gluon incoming momentum q and colour polarization e^a , the second equalities are valid only for an infinitesimal q (or in the case of an abelian external field), and $(t^a)^{bc} = if^{acb}$ in the adjoint representation. Therefore, the QCD proper vertex $\Lambda_\mu^a(p, q) = \Lambda_\mu t^a$ obeys $\Lambda_\mu^a q^\mu e^a = -\Sigma(p+qe^{at^a}) + \Sigma(p)$ for infinitesimal q , or $\Lambda_\mu(p, 0) = -\frac{\partial \Sigma(p)}{\partial p^\mu}$. The form factor is projected out by $\varepsilon(0) = Z_Q [1 + \frac{1}{4} \text{Tr} \Lambda_\mu v^\mu (1 + \not{v})]$. On the mass shell, $\frac{1}{4} \text{Tr} \frac{\partial \Sigma}{\partial p^\mu} = (1 - Z_Q^{-1}) v_\mu$, and hence $\varepsilon(0) = 1$.

Let's denote the sum of bare vertex diagrams in HQET at $1/m^0$ as $igt^a v^\mu \frac{1+\not{v}}{2} [1 + \Lambda(\omega, \Delta)]$, where $\Delta = qv = \omega' - \omega$. The Ward identity for the static quark propagator is the same as for the ordinary one (Fig. 2). Therefore, $\Delta e^{at^a} \Lambda(\omega, \Delta) = -\Sigma(\omega + \Delta e^{at^a}) + \Sigma(\omega)$ for infinitesimal Δ , or

$$\Lambda(\omega, 0) = -\frac{d\Sigma(\omega)}{d\omega}. \quad (13)$$

It is interesting, that for an abelian external field $\Lambda(\omega, \Delta) = -\frac{\Sigma(\omega+\Delta) - \Sigma(\omega)}{\Delta}$ exactly. The total colour charge of a static quark $Z_Q [1 + \Lambda(0, 0)] = 1$, as expected.

The $1/m$ HQET bare proper vertex has the form

$$i\frac{C_k}{2m} gt^a \frac{1+\not{v}}{2} \left[(1 + \Lambda_k)(p+p')_{\perp}^\mu + (\Lambda_{k0} + \Lambda_{k1} p_{\perp}^2 + \Lambda'_{k1} p_{\perp}'^2 + \Lambda_{k2} q_{\perp}^2) v^\mu \right] + i\frac{C_m}{4m} gt^a \frac{1+\not{v}}{2} [\gamma^\mu, \not{v}] \frac{1+\not{v}}{2} (1 + \Lambda_m), \quad (14)$$

where all Λ_i depend on ω, Δ ; $\Lambda'_{k1}(\omega, \Delta) = \Lambda_{k1}(\omega + \Delta, -\Delta)$; $\Lambda_k(\omega, \Delta) = \Lambda_k(\omega + \Delta, -\Delta)$, and similarly for $\Lambda_{k0}, \Lambda_{k2}$. Similarly to the previous Section, we can see that variation of the leading vertex function at $v \rightarrow v + \delta v$ coincides with that of the kinetic-energy vertex function at $p_{\perp} \rightarrow p_{\perp} + \delta p_{\perp}$, if $\delta v = \frac{C_k}{m} \delta p_{\perp}$. This requires

$$\Lambda_k(\omega, \Delta) = \Lambda(\omega, \Delta), \quad \Lambda'_{k1}(\omega, \Delta) = \frac{\partial \Lambda(\omega, \Delta)}{\partial \Delta} \quad (15)$$

(and hence $\Lambda_{k1}(\omega, \Delta) = (\frac{\partial}{\partial \omega} - \frac{\partial}{\partial \Delta}) \Lambda(\omega, \Delta)$). The Ward identities of Fig. 2 result in

$$\Lambda_{k0}(\omega, 0) = -\frac{d\Sigma_{k0}(\omega)}{d\omega}, \quad \Lambda_{k2}(\omega, 0) = 0 \quad (16)$$

(in an abelian external field, $\Lambda_{k0}(\omega, \Delta) = -\frac{\Sigma_{k0}(\omega+\Delta) - \Sigma_{k0}(\omega)}{\Delta}$, $\Lambda_{k2}(\omega, \Delta) = 0$).




Figure 2: Ward identities in the background field formalism

Reparameterization invariance relates the spin-orbit vertex function to the chromomagnetic one, but we shall not discuss details here.

The on-shell HQET vertex at the tree level is

$$\bar{u}_v(k') \left(v^\mu + C_k \frac{(k+k')^\mu}{2m} + C_m \frac{[\not{q}, \gamma^\mu]}{4m} + C_d \frac{q^2}{8m^2} v^\mu + C_s \frac{[\not{k}, \not{q}]}{8m^2} v^\mu + \dots \right) u_v(k) \quad (17)$$

As we have demonstrated above, there are no corrections to the first two terms. Other terms have corrections starting from two loops, if there is a finite-mass flavour (such as c in b -quark HQET). Expressing the on-shell QCD vertex via HQET spinors, we obtain

$$\bar{u}_v(k') \left[\varepsilon(q^2) \left(v^\mu + \frac{(k+k')^\mu}{2m} - \frac{q^2 + [\not{k}, \not{q}]}{8m^2} v^\mu + \dots \right) + \mu(q^2) \left(\frac{[\not{q}, \gamma^\mu]}{4m} + \frac{q^2 + [\not{k}, \not{q}]}{4m^2} v^\mu + \dots \right) \right] u_v(k). \quad (18)$$

Therefore, the coefficients in the HQET lagrangian are

$$C_k = 1, \quad C_m = \mu, \quad C_d = 8\varepsilon' + 2\mu - 1, \quad C_s = 2\mu - 1. \quad (19)$$

The first one has no corrections (10). The coefficients (19) are not independent:

$$C_s = 2C_m - 1. \quad (20)$$

Probably, reparameterization-invariance Ward identities yield relations among corrections from finite-mass loops in HQET which ensure the absence of corrections to (20). However, we shall not trace details here.

Similarly, at the $1/m^3$ level, the coefficients in the HQET lagrangian are

$$C_{w1} = 4\mu' + \frac{1}{2}\mu + \frac{1}{2}, \quad C_{w2} = 4\mu' + \frac{1}{2}\mu - \frac{1}{2}, \quad C_{p'p} = \mu - 1, \quad C_M = -4\varepsilon' - \frac{1}{2}\mu + \frac{1}{2} \quad (21)$$

They are not independent:

$$C_{w2} = C_{w1} - 1, \quad C_{p'p} = C_m - 1, \quad C_M = \frac{1}{2}(C_m - C_d). \quad (22)$$

Calculation of C_a, C_b requires matching amplitudes with two gluons. Calculation of contact terms requires matching amplitudes with light quarks.

4. Chromomagnetic interaction at two loops

As we know, the kinetic coefficient $C_k(\mu) = 1$, and the only coefficient in the HQET lagrangian up to $1/m$ level which is not known exactly is the chromomagnetic coefficient $C_m(\mu)$. It is natural to find it from QCD/HQET matching at $\mu \sim m$ where no large logarithms appear. Renormalization group can be used to obtain C_m at $\mu \ll m$:

$$C_m(\mu) = C_m(m) \exp \left(- \int_{\alpha_s(m)}^{\alpha_s(\mu)} \frac{\gamma_m(\alpha) d\alpha}{2\beta(\alpha) \alpha} \right), \quad (23)$$

where $C_m(m) = 1 + C_1 \frac{\alpha_s(m)}{4\pi} + C_2 \left(\frac{\alpha_s(m)}{4\pi} \right)^2 + \dots$, $\gamma_m = \frac{d \log Z_m}{d \log \mu} = \gamma_1 \frac{\alpha_s}{4\pi} + \gamma_2 \left(\frac{\alpha_s}{4\pi} \right)^2 + \dots$ is the anomalous dimension of the chromomagnetic operator in HQET, and the β -function is $\beta = -\frac{1}{2} \frac{d \log \alpha_s}{d \log \mu} = \beta_1 \frac{\alpha_s}{4\pi} + \beta_2 \left(\frac{\alpha_s}{4\pi} \right)^2 + \dots$ (where $\beta_1 = \frac{11}{3} C_A - \frac{4}{3} T_F n_f$). If $L = \log m/\mu$ is not very large, it is better to retain all two-loop terms and neglect higher loops:

$$C_m(\mu) = 1 + (C_1 - \gamma_1 L) \frac{\alpha_s(m)}{4\pi} + [C_2 - (C_1 \gamma_1 + \gamma_2) L + \gamma_1 (\gamma_1 - \beta_1) L^2] \left(\frac{\alpha_s}{4\pi} \right)^2 \quad (24)$$

This approximation holds up to relatively large L because C_2 is numerically large. If L is parametrically large, then it is better to sum leading and subleading logarithms:

$$C_m(\mu) = \left(\frac{\alpha_s(\mu)}{\alpha_s(m)} \right)^{-\frac{\gamma_1}{2\beta_1}} \left[1 + C_1 \frac{\alpha_s(m)}{4\pi} - \frac{\beta_1 \gamma_2 - \beta_2 \gamma_1}{2\beta_1^2} \frac{\alpha_s(\mu) - \alpha_s(m)}{4\pi} \right]. \quad (25)$$

In this case, we cannot utilize C_2 without knowing γ_3 . In general, the solution of (23) can be written as

$$C_m(\mu) = \hat{C}_m K(\mu), \quad \hat{C}_m = \alpha_s(m)^{\frac{\gamma_1}{2\beta_1}} (1 + \delta c), \quad \delta c = c_1 \frac{\alpha_s(m)}{4\pi} + c_2 \left(\frac{\alpha_s(m)}{4\pi} \right)^2 + \dots \quad (26)$$

where \hat{C}_m is scale- and scheme-independent.

As a simple application, we consider $B-B^*$ mass splitting [28,29]¹

$$m_{B^*} - m_B = \frac{2C_m(\mu)}{3m} \mu_m^2(\mu) \quad (27)$$

¹in [28], ρ_{mm}^3 is missing; in [29], the leading logarithmic running of $C_m(\mu)$ has a wrong sign.

$$+ \frac{1}{3m^2} [C_m(\mu) \rho_{km}^3(\mu) + C_m^2(\mu) \rho_{mm}^3(\mu) - C_s(\mu) \rho_s^3(\mu)],$$


where $\mu_m^2(\mu)$ and $\rho_s^3(\mu)$ are local matrix elements of chromomagnetic interaction and spin-orbit one, while $\rho_{km}^3(\mu)$ and $\rho_{mm}^3(\mu)$ are kinetic-chromomagnetic and chromomagnetic-chromomagnetic bilocal matrix elements (in the later case, there are two γ -matrix structures, 1 and $\sigma_{\mu\nu}$; the coefficient of the second one is implied here). Introducing renormalization group invariants

$$\hat{\mu}_m^2 = K(\mu) \mu_m^2(\mu), \quad \hat{\rho}_{km}^3 = K(\mu) \rho_{km}^3(\mu) + [1 - K(\mu)] \rho_s^3(\mu), \quad \hat{\rho}_{mm}^3 = K^2(\mu) \rho_{mm}^3(\mu), \quad \hat{\rho}_s^3 = \rho_s^3(\mu), \quad (28)$$

we can rewrite it as

$$m_{B^*} - m_B = \frac{2\hat{C}_m}{3m} \hat{\mu}_m^2 + \frac{1}{3m^2} [\hat{C}_m (\hat{\rho}_{km}^3 - 2\hat{\rho}_s^3) + \hat{C}_m^2 \hat{\rho}_{mm}^3 + \hat{\rho}_s^3]. \quad (29)$$

In order to obtain C_m , we should calculate the heavy-quark chromomagnetic moment μ (Fig. 3). All on-shell massive integrals can be reduced to 3 basis ones



$$I_0^2 = \text{diagram}, \quad I_1 = \text{diagram}, \quad I_2 = \text{diagram} \quad (30)$$

using integration by parts [30]–[32]. I_0^2 and I_1 are expressed via Γ -functions of d ; I_2 is expressed via I_0^2 , I_1 , and one difficult convergent integral [32]

$$I = \pi^2 \log 2 - \frac{3}{2} \zeta(3) + O(\epsilon). \quad (31)$$

The result has the structure

$$\mu = 1 + \frac{g_0^2 m^{-2\epsilon}}{(4\pi)^{d/2}} (C_F, C_A) \times I_0 + \frac{g_0^4 m^{-4\epsilon}}{(4\pi)^d} \times (C_F^2, C_F C_A, C_A^2, C_F T_F n_f, C_A T_F n_f, C_F T_F, C_A T_F) \times (I_0^2, I_1, I_2). \quad (32)$$

Now we express it via $\alpha_s(\mu)$ and expand in ϵ . The coefficient of $1/\epsilon$ gives the anomalous dimension

$$\gamma_m = 2C_A \frac{\alpha_s}{4\pi} + \frac{4}{9} C_A (17C_A - 13T_F n_f) \left(\frac{\alpha_s}{4\pi} \right)^2 + \dots \quad (33)$$




Figure 3: Diagrams for the QCD proper vertex

The chromomagnetic interaction coefficient at $\mu = m$ is

$$\begin{aligned}
 (36) \quad C_m(m) &= 1 + 2(C_F + C_A) \frac{\alpha_s(m)}{4\pi} \\
 &+ \left[C_F^2 \left(-8I + \frac{20}{3}\pi^2 - 31 \right) + C_F C_A \left(\frac{4}{3}I + \frac{4}{3}\pi^2 + \frac{269}{9} \right) \right. \\
 &+ C_A^2 \left(\frac{4}{3}I - \frac{17}{9}\pi^2 + \frac{805}{27} \right) \\
 &+ C_F T_F n_l \left(-\frac{100}{9} \right) + C_A T_F n_l \left(-\frac{4}{9}\pi^2 - \frac{299}{27} \right) \\
 &\left. + C_F T_F \left(-\frac{16}{3}\pi^2 + \frac{476}{9} \right) + C_A T_F \left(\pi^2 - \frac{298}{27} \right) \right] \left(\frac{\alpha_s}{4\pi} \right)^2 \\
 &= 1 + \frac{13}{6} \frac{\alpha_s(m)}{\pi} + (21.79 - 1.91n_l) \left(\frac{\alpha_s}{\pi} \right)^2.
 \end{aligned} \quad (34)$$

The coefficient of $(\alpha_s/\pi)^2$ is about 11 for $n_l = 4$ light flavours. It is 40% less than the expectation based on naive nonabelianization [33]. The contribution of the heavy quark loop to this coefficient is merely -0.1 .

5. Chromomagnetic interaction at higher loops

Perturbation series for C_m can be rewritten via β_1 instead of n_f :

$$C_m(\mu) = 1 + \sum_{L=1}^{\infty} \sum_{n=0}^{L-1} a_{Ln} \beta_1^n \alpha_s^L = 1 + \frac{1}{\beta_1} f(\beta_1 \alpha_s) + O\left(\frac{1}{\beta_1^2}\right). \quad (35)$$

There is no sensible limit of QCD in which β_1 may be considered a large parameter (except, may be, $n_f \rightarrow -\infty$). However, retaining only the leading β_1 terms often gives a good approximation to exact multi-loop results [33]. This limit is believed to provide information about summability of perturbation series [34]. At the first order in $1/\beta_1$, multiplicative renormalization amounts to subtraction of $1/\epsilon^n$ terms;

$$\frac{\beta_1 g_0^2}{(4\pi)^2} = \bar{\mu}^{2\epsilon} \frac{\beta}{1 + \beta/\epsilon}, \quad \beta = \frac{\beta_1 \alpha_s}{4\pi} = \frac{1}{2 \log \mu / \Lambda_{\overline{\text{MS}}}}. \quad (36)$$

The perturbation series (35) can be rewritten as

$$C_m(\mu) = 1 + \frac{1}{\beta_1} \sum_{L=1}^{\infty} \frac{F(\epsilon, L\epsilon)}{L} \left(\frac{\beta}{\epsilon + \beta} \right)^L - (\text{subtractions}) + O\left(\frac{1}{\beta_1^2}\right). \quad (37)$$

Knowledge of the function $F(\varepsilon, u)$ allows one to obtain the anomalous dimension

$$\gamma_m = \frac{2\beta}{\beta_1} F(-\beta, 0) + O\left(\frac{1}{\beta_1^2}\right) \quad (38)$$

and the finite term

$$C_m(\mu) = 1 + \frac{1}{\beta_1} \int_{-\beta}^0 d\varepsilon \frac{F(\varepsilon, 0) - F(0, 0)}{\varepsilon} + \frac{1}{\beta_1} \int_0^\infty du e^{-u/\beta} \frac{F(0, u) - F(0, 0)}{u} + O\left(\frac{1}{\beta_1^2}\right) \quad (39)$$

(this method was used in [33]; see references in this paper). Renormalization group invariant (26) is

$$\delta c = \frac{1}{\beta_1} \int_0^\infty du e^{-\frac{4\pi}{\beta_1 \alpha_s} u} S(u) + O\left(\frac{1}{\beta_1^2}\right), \quad (40)$$

$$S(u) = e^{-\frac{5}{3}u} \frac{F(0, u) - F(0, 0)}{u} \Big|_{\mu=m}$$

(here α_s is taken at $\mu = m$ in the V -scheme, $\exp(-\frac{4\pi}{\beta_1 \alpha_s} u) = (\frac{\Lambda_V}{m})^{-2u}$).




Figure 4: L -loop diagrams with the maximum number of quark loops.

The function $F(\varepsilon, u)$ is determined by the coefficient of the highest degree of n_f in the L -loop term, which is given by the diagrams in Fig. 4. Calculating them, we obtain

$$F(\varepsilon, u) = \left(\frac{\mu}{m}\right)^{2u} e^{\gamma\varepsilon} \frac{\Gamma(1+u)\Gamma(1-2u)}{\Gamma(3-u-\varepsilon)} D(\varepsilon)^{u/\varepsilon-1} N(\varepsilon, u)$$

$$D(\varepsilon) = 6e^{\gamma\varepsilon} \Gamma(1-\varepsilon) B(2-\varepsilon, 2-\varepsilon) = 1 + \frac{5}{3}\varepsilon + \dots \quad (41)$$

$$N(\varepsilon, u) = C_F 4u(1+u-2\varepsilon u) + C_A \frac{2-u-\varepsilon}{2(1-\varepsilon)} (2+3u-5\varepsilon-6\varepsilon u+2\varepsilon^2+4\varepsilon^2 u).$$

This gives the anomalous dimension

$$\gamma_m = C_A \frac{\alpha_s}{2\pi} \frac{\beta(1+2\beta)\Gamma(5+2\beta)}{24(1+\beta)\Gamma^3(2+\beta)\Gamma(1-\beta)} = C_A \frac{\alpha_s}{2\pi} \left[1 + \frac{13}{6} \frac{\beta_1 \alpha_s}{4\pi} - \frac{1}{2} \left(\frac{\beta_1 \alpha_s}{4\pi} \right)^2 + \dots \right]. \quad (42)$$

This perturbation series is convergent with the radius $\beta_1 |\alpha_s| < 4\pi$. The Borel image of δc

$$S(u) = \frac{\Gamma(u)\Gamma(1-2u)}{\Gamma(3-u)} [4u(1+u)C_F + \frac{1}{2}(2-u)(2+3u)C_A] - e^{-\frac{5}{3}u} \frac{C_A}{u} \quad (43)$$

has infrared renormalon poles at $u = \frac{n}{2}$. They produce ambiguities in the sum of the perturbation series for δc , which are of order of the residues $\sim (\Lambda_V/m)^n$. The leading ambiguity ($u = \frac{1}{2}$) is

$$\Delta \hat{C}_m = \left(1 + \frac{7}{8} \frac{C_A}{C_F} \right) \frac{\Delta m}{m}, \quad (44)$$

where Δm is the ambiguity of the heavy-quark pole mass [35,36].

Physical quantities, such as the mass splitting (27), are factorized into short-distance coefficients and long-distance hadronic matrix elements. In regularization schemes without a hard momentum cut-off, such as $\overline{\text{MS}}$, Wilson coefficients also contain large-distance contributions which produce infrared renormalon ambiguities. Likewise, hadronic matrix elements contain small-distance contributions which produce ultraviolet renormalon ambiguities. In other words, the separation into short- and long-distance contributions is ambiguous; only when they are combined to form a physical quantity, an unambiguous result is obtained. Cancellations between infrared and ultraviolet renormalon ambiguities in HQET were traced in [37].

Ultraviolet renormalon ambiguities in matrix elements ρ_i^3 don't depend on external states, and may be calculated at the level of quarks and gluons (Fig. 5). Note that there is an ultraviolet renormalon ambiguity in the wave function renormalization $\Delta Z_Q = \frac{3}{2} \frac{\Delta m}{m}$ (Fig. 5d). The result is

$$\Delta \rho_{km}^3 = -\frac{2}{3} \frac{C_A}{C_F} \mu_m^2 \Delta m, \quad \Delta \rho_{mm}^3 = -\frac{19}{12} \frac{C_A}{C_F} \mu_m^2 \Delta m, \quad \Delta \rho_s^3 = -\frac{1}{2} \frac{C_A}{C_F} \mu_m^2 \Delta m \quad (45)$$




Figure 5: Diagrams for ρ_i^3 ; quark loops are inserted in all possible ways.

The sum of ultraviolet ambiguities of the $1/m^2$ contributions to (27) cancels the infrared ambiguity of the leading term.

The requirement of cancellation of renormalon ambiguities in the mass splitting (28) for all m allows us to establish the structure of the leading infrared renormalon singularity in $S(u)$ at $u = \frac{1}{2}$ beyond the large β_1 limit. The ultraviolet ambiguity of the square bracket in (28) should be equal to $\bar{\mu}_m^2$ times

$$\Lambda_V = m e^{-\frac{2\pi}{\beta_1 \alpha_s}} \alpha_s^{-\frac{\beta_2}{2\beta_1^2}} [1 + O(\alpha_s)]. \quad (46)$$

In order to reproduce the correct fractional powers of α_s , $S(u)$ in (40) should

have the branch point at $u = \frac{1}{2}$ instead of a pole:

$$S(u) = \frac{1}{(\frac{1}{2} - u)^{1+\beta_2/2\beta_1^2}} \left[2C_F K_1 - \frac{1}{3} C_A K_2 \right. \\ \left. + \frac{19}{12} \frac{C_A K_3}{(\frac{1}{2} - u)^{-\gamma_1/2\beta_1}} + \frac{1}{2} \frac{C_A K_4}{(\frac{1}{2} - u)^{\gamma_1/2\beta_1}} \right] \quad (47)$$

where omitted terms are suppressed as $\frac{1}{2} - u$ compared to the displayed ones. Normalization constants are known in the large β_1 limit only: $K_i = 1 + O(1/\beta_1)$. The large-order behaviour of the perturbation series for δc is

$$c_{n+1} = n! (2\beta_1)^n n^{\beta_2/2\beta_1^2} \left[4C_F K_1 - \frac{2}{3} C_A K_2 \right. \\ \left. + \frac{19}{6} C_A K_3 n^{-\gamma_1/2\beta_1} + C_A K_4 n^{\gamma_1/2\beta_1} \right] \quad (48)$$

where omitted terms are suppressed as $1/n$ compared to the displayed ones.

Acknowledgements. I am grateful to A. Czarnecki and M. Neubert for collaboration in writing [19,20]; to S. Groote for ongoing collaboration; to C. Balzereit for discussing [10,26]; to T. Mannel for useful discussions; to J. G. Körner for hospitality at Mainz during preparation of this talk; and to M. Beyer for organization of the workshop.

References

- [1] M. Neubert, Phys. Reports **245** (1994) 259.
- [2] E. Eichten and B. Hill, Phys. Lett. **B234** (1990) 511.
- [3] E. Eichten and B. Hill, Phys. Lett. **B243** (1990) 427.
- [4] A. F. Falk, B. Grinstein, and M. E. Luke, Nucl. Phys. **B357** (1991) 185.
- [5] C. L. Y. Lee, Preprint CALT-68-1663 (1991); revised (1997).
- [6] A. V. Manohar, Phys. Rev. **D56** (1997) 230.
- [7] M. Luke and A. V. Manohar, Phys. Lett. **B286** (1992) 348.
- [8] Y.-Q. Chen, Phys. Lett. **B317** (1993) 421.
- [9] W. Kilian and T. Ohl, Phys. Rev. **D50** (1994) 4649.
- [10] C. Balzereit, Diploma thesis, Darmstadt (1994).
- [11] M. Finkemeier, H. Georgi, and M. McIrvin, Phys. Rev. **D55** (1997) 6933.
- [12] R. Sundrum, Preprint BUHEP-97-14, hep-ph/9704256 (1997); Phys. Rev. **D57**, in print.

- [13] C. L. Y. Lee, Preprint UCSD-TH-97-24, hep-ph/9709238 (1997).
 [14] J. G. Körner and G. Thompson, Phys. Lett. **B264** (1991) 185.
 [15] S. Balk, J. G. Körner, and D. Pirjol, Nucl. Phys. **B428** (1994) 499.
 [16] T. Mannel, W. Roberts, and Z. Ryzak, Nucl. Phys. **B368** (19204) 92.
 [17] C. L. Y. Lee, Preprint UCSD-TH-97-23, hep-ph/9709237 (1997).
 [18] G. Amorós, M. Beneke, and M. Neubert, Phys. Lett. **B401** (1997) 81.
 [19] A. Czarnecki and A. G. Grozin, Phys. Lett. **B405** (1997) 142.
 [20] A. G. Grozin and M. Neubert, Preprint CERN-TH/97-102, hep-ph/9707318 (1997); Nucl. Phys. **B**, in print.
 [21] Y.-Q. Chen, Y. P. Kuang, and R. Oakes, Phys. Rev. **D52** (1995) 264.
 [22] C. Balzereit and T. Ohl, Phys. Lett. **B386** (1996) 335.
 [23] M. Finkemeier and M. McIrvin, Phys. Rev. **D55** (1997) 377.
 [24] B. Blok, J. G. Körner, D. Pirjol, and J. C. Rojas, Nucl. Phys. **B496** (1997) 358.
 [25] C. Bauer and A. V. Manohar, Preprint UCSD/PTH 97-19, UTP-97-17, hep-ph/9708306 (1997); Phys. Rev. **D57**, in print.
 [26] C. Balzereit, These Proceedings.
 [27] L. F. Abbott, Acta Phys. Polonica **13** (1982) 33.
 [28] T. Mannel, Phys. Rev. **D50** (1994) 428.
 [29] I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D52** (1995) 196.
 [30] N. Gray, D. J. Broadhurst, W. Grafe, and K. Schilcher, Zeit. Phys. **C48** (1990) 673.
 [31] D. J. Broadhurst, N. Gray, and K. Schilcher, Zeit. Phys. **C52** (1991) 111.
 [32] D. J. Broadhurst, Zeit. Phys. **C54** (1992) 599.
 [33] D. J. Broadhurst and A. G. Grozin, Phys. Rev. **D52** (1995) 4082.
 [34] A. H. Mueller, QCD 20 years later, ed. P. M. Zerwas and H. A. Kastrup, World Scientific (1993), p. 162.
 [35] M. Beneke and V. M. Braun, Nucl. Phys. **B426** (1994) 301.
 [36] I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D50** (1994) 2234.
 [37] M. Neubert and C. T. Sachrajda, Nucl. Phys. **B438** (1995) 235.

A.G. Grozin

Recent progress on HQET lagrangian

А.Г. Грозин

**Недавний прогресс
касательно лагранжиана HQET**

Budker INP 97-90

Ответственный за выпуск А.М. Кудрявцев

Работа поступила 29.10. 1997 г.

Сдано в набор 12.11.1997 г.

Подписано в печать 13.11.1997 г.

Формат бумаги 60×90 1/16 Объем 0.9 печ.л., 0.8 уч.-изд.л.

Тираж 100 экз. Бесплатно. Заказ № 90

Обработано на IBM PC и отпечатано на

ротапинтере ИЯФ им. Г.И. Будкера СО РАН,

Новосибирск, 630090, пр. академика Лаврентьева, 11.